

1 Title: The influence of early exercise postpartum on pelvic floor muscle function and prevalence  
2 of pelvic floor dysfunction 12 months postpartum.

3 Authors

4 1. Merete Kolberg Tennfjord <sup>1,2,3</sup>

5 2. Marie Ellström Engh <sup>2,4</sup>

6 3. Kari Bø <sup>1,2</sup>

7

8 <sup>1</sup> Norwegian School of Sport Sciences, Department of Sports Medicine, Oslo, Norway.

9 <sup>2</sup> Akershus University Hospital, Department of Obstetrics and Gynaecology, Lørenskog, Norway.

10 <sup>3</sup> Kristiania University College, School of Health Science, Oslo, Norway.

11 <sup>4</sup> Institute of Clinical Medicine, University of Oslo, Oslo, Norway

12

13 Full correspondence details

Name	Assoc Prof. Merete Kolberg Tennfjord
Country	Norway
Mob	+4798663431
Email	merete.kolberg@gmail.com

14

15

16

17

18 ABSTRACT

19 Background: There is limited knowledge on how exercise impacts on the pelvic floor muscles  
20 (PFM) and prevalence of stress urinary incontinence (SUI) and pelvic organ prolapse (POP)  
21 postpartum.

22 Objective: To investigate if early onset of general exercise postpartum negatively affects the PFM  
23 and/or increases the risk of SUI and POP 12 months postpartum?

24 Design: Prospective cohort study

25 Methods: At 6 weeks postpartum, 57 women classified as Exercisers ( $\geq 3$  times  $\geq 30$  min/week)  
26 were compared with 120 Non-exercisers (mean age 29 years, SD 4.3). Manometry was used to  
27 measure vaginal resting pressure (VRP), PFM strength and PFM endurance, and symptoms of  
28 SUI and POP were assessed by questionnaires. Data were presented as standardised beta  
29 coefficients (B) and odds ratio (OR).

30 Results: No differences were found between Exercisers (n=57) and Non-exercisers (n=120) at 6  
31 weeks postpartum on VRP: B -0.04 (95%CI -3.4, 2.1), PFM strength: B 0.03 (95%CI -4.7, 7.4),  
32 PFM endurance: B -0.02 (95%CI -59, 46), or symptoms of SUI: OR 0.51 (95%CI: 0.25, 1.1) or  
33 POP: OR 0.62 (95%CI: 0.26, 1.5) measured at 12 months postpartum. Adjusting for covariates,  
34 women with BMI between 25-29.9 and BMI  $>30$  were more likely to report SUI 12 months  
35 postpartum (OR=2.2, 95%CI: 1.0, 4.7 and OR=3.3, 95%CI: 1.2, 9.4, respectively). Women with  
36 physically strenuous occupations were more likely to report POP 12 months postpartum  
37 (OR=3.0, 95%CI: 1.2, 7.3).

38 Limitations: No sample size calculation was undertaken for this study.

39 Conclusion: This study suggests that regular exercise 6 weeks postpartum has no negative effect  
40 on PFM function or SUI or POP. Being overweight, however, was associated with more SUI, and  
41 women with physically strenuous occupations reported more POP.

42

43 Manuscript word count: 3340

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 INTRODUCTION

60 Guidelines regarding exercise for postpartum women recommend a gradual commencement or  
61 return to physical activity as soon as medically safe (1). Concerns have been raised, however, that  
62 starting exercise early in the postpartum period may negatively impact on the already stretched  
63 and weakened pelvic floor muscles (PFM) (2-4). Transperineal ultrasound has shown changes in  
64 pelvic organ support and an increase in the area surrounding the pelvic openings (levator hiatus  
65 area) during both pregnancy and following vaginal delivery, which suggests reduced support to  
66 the pelvic organs (3,4). These changes are still present at 6 weeks and 12 months postpartum (3).  
67 Both pregnancy and childbirth may increase the risk of developing pelvic floor dysfunctions  
68 (PFD) such as urinary incontinence (UI), pelvic organ prolapse (POP) and anal incontinence (AI)  
69 (5). At 12 months postpartum, prevalence rates of up to 30% for UI (most commonly stress UI  
70 (SUI)), and 19% for both POP and AI have been found (6-8).

71 In 2004, Bø et al (9) described two contrasting hypotheses regarding general exercise and its  
72 effect on the pelvic floor: 1. General exercise strengthens the PFM and 2. General exercise  
73 weakens the PFM. To date there is limited knowledge on the effects of commencing general  
74 exercise early in the postpartum period on the PFM and the risks of developing SUI, POP and AI  
75 (10).

76 The research questions of the present study were:

- 77 1) Does general exercise commenced early in the postpartum period affect PFM strength and  
78 endurance 12 months following delivery?
- 79 2) Does general exercise commenced early in the postpartum period have an effect on  
80 symptoms of SUI, POP and AI 12 months following delivery?

## 81 METHODS

### 82 Design

83 All nulliparous women scheduled for delivery at Akershus University Hospital between January  
84 2010 and April 2011 were invited to participate in this cohort study. Three hundred women of  
85 mean gestational week 20.9 (SD 1.4) were recruited. Questionnaires regarding PFD were  
86 administered at 6 different stages: pre-pregnancy; gestational week 22 and 37; and 6 weeks, 6  
87 months and 12 months postpartum. Questionnaires for the pre-pregnancy stage were administered  
88 retrospectively. The present study used only questionnaire data at 6 weeks (baseline) and 12  
89 months postpartum. Vaginal resting pressure (VRP), PFM strength and PFM endurance were  
90 recorded at 6 weeks (baseline) and 12 months postpartum. Relationships were analysed between  
91 Exercises and Non-exercises on these variables at both points in time as were symptoms of SUI,  
92 POP and AI. This study followed the recommended STROBE reporting guidelines:  
93 *Strengthening the reporting of observational studies in epidemiology* (11).

### 94 Participants

95 This study recruited primiparous women who had given birth to a singleton baby and who were  
96 able to understand a Scandinavian language. Exclusion criteria were: multiple pregnancies, past  
97 history of miscarriage after gestational week 16, preterm delivery (<week 32 gestation), stillbirth,  
98 serious illness to mother or child, a subsequent pregnancy of 6 weeks gestation or more , and  
99 recruitment to a parallel project investigating effects of PFM training (PFMT) (ClinicalTrials.gov  
100 NCT01069484). The Regional Medical Ethics Committee (2009/170) (approval date 04.08.09)  
101 and the Norwegian Social Science Data Services (2799026) (approval date 28.09.09) approved  
102 this study. All participants gave written informed consent to participate prior to entering the

103 study.

#### 104 Exercise

105 Exercise is defined as: "Physical activity that is usually performed on a repeated basis over an  
106 extended period of time with a specific aim, such as to improve fitness, physical performance or  
107 health" (12). In the questionnaires, participants reported how often they performed the following  
108 types of exercise: strolling, brisk walking, running (jogging or orienteering), bicycling, skiing,  
109 swimming, ball games, weight training , low-impact aerobics classes, high-impact aerobics  
110 classes, special fitness classes for pregnant and postpartum women, dancing (swing, rock and roll  
111 and folk), horseback riding and other forms of exercise. For the purposes of this study, strolling  
112 was not classified as exercise (13). The weekly exercise frequency score was summed across all  
113 13 exercise categories. Participants were classified as Exercises if they trained "≥3 times ≥30  
114 min/week" at 6 weeks postpartum (14). Those who exercised 0-2 times per week were considered  
115 Non-exercisers.

#### 116 Outcome measures

117 *Primary outcomes:* VRP, PFM strength and PFM endurance were measured using a high  
118 precision pressure transducer connected to an air-filled vaginal balloon (Camtech AS, Sandvika,  
119 Norway). Before measurements were taken, participants were given a short anatomy lecture and  
120 taught how to correctly contract their PFM using observation and vaginal palpation (15). The  
121 pressure transducer was positioned such that the middle of the balloon was inserted into the  
122 vagina 3.5cm from the introitus where the vaginal high pressure zone is located (16). This  
123 measurement method has demonstrated good validity and reliability with simultaneous  
124 observation of an inward movement of the catheter and no use of muscles of the abdomino-pelvic  
125 cavity (17-19). VRP (cmH<sub>2</sub>O) was calculated as the difference between the atmospheric pressure

126 and the vaginal high pressure zone at rest. PFM strength (cmH<sub>2</sub>O) was measured from the resting  
127 pressure line to the peak, not including the resting pressure, and reported as the mean of 3  
128 maximal voluntary contractions. PFM endurance (cmH<sub>2</sub>Osec) was assessed as the area under the  
129 curve of a single 10 second maximal contraction (20). All measurements were taken with the  
130 participant in hook lying with one leg resting against a wall. Two female physiotherapists (with  
131 specialization in Women's Health) performed the measurements. These therapists had  
132 demonstrated good intra-rater and inter-rater reliability (Intraclass correlation coefficients >0.91)  
133 for all test measurements (VRP, PFM strength and PFM endurance) prior to the commencement  
134 of the study (non-published data).

135 *Secondary outcomes:* Prevalence of symptoms of SUI, POP and AI were measured using the  
136 International Consultation on Incontinence Modular Questionnaire (ICIQ). For SUI, the validated  
137 Norwegian version of the ICIQ – Urinary Incontinence short form (ICIQ-UI Short Form) was  
138 used (21,22). Women were classified with SUI if they answered that they leaked when they:  
139 “cough or sneeze”, or when “physically active/exercising” to the question: “When do you leak  
140 urine”?

141 For symptoms of POP, the ICIQ – vaginal symptoms questionnaire (ICIQ-VS) (23) was used.  
142 Participants were classified as not having POP if they answered “never” to the questions: “Are  
143 you aware of a lump or bulge in your vagina?” and “Can you feel or see a lump or a bulge on the  
144 outside of your vagina?” Responses to both these questions were pooled into one category for the  
145 purpose of analyses. The ICIQ-VS was translated into Norwegian by the study group during the  
146 planning of the study but as yet has not undergone linguistic validation.

147 For AI, the ICIQ – bowel symptoms questionnaire (ICIQ-B) (24) was used. Participants were  
148 classified as incontinent, based on the definitions from Sultan et al (25), if they answered “never”

149 to the questions: “Are you able to control watery or loose stools from your back passage?”, “Are  
150 you able to control accidental loss of formed or solid stools from your back passage?” and “Are  
151 you able to control wind (flatus) escaping from your back passage?” The ICIQ-B was translated  
152 into Norwegian by the study group and has not yet undergone linguistic validation.

### 153 Data analysis

154 Background variables for the Exercisers and Non-exercisers were measured at 6 weeks  
155 postpartum and presented as frequencies (n) and percentages, or means with standard deviations  
156 (SD). Continuous variables were normally distributed as assessed by a Normal Q-Q plot.

157 To assess differences between Exercisers and Non-exercisers at 6 weeks postpartum on VRP,  
158 PFM strength and PFM endurance at 12 months postpartum independent sample t-test were used.

159 Differences between to the two groups on symptoms of SUI, POP and AI were assessed at 12  
160 months postpartum using the Chi-Square test. Possible outliers were assessed by Q-Q plot. Due  
161 to only two women (1.1%) reporting symptoms of AI at 12 months postpartum, AI was not  
162 included in any further analyses. Standard multiple linear regression and logistic regression was  
163 used to adjust for possible covariates. Potential covariates based on previously published  
164 literature, and clinical reasoning on risk factors for reduced PFM function and presence of SUI  
165 and POP (26), were assessed for eligibility and included in the regression models. Covariates  
166 included in our study were: physically strenuous occupation (yes/no), BMI, age, performing  
167 PFMT  $\geq$  three times per week, and the use of vacuum/forceps (yes/no). Questions asked  
168 regarding occupation included: “Do you consider your work physically demanding?” The  
169 answers were pooled into yes (for those who responded both yes and sometimes)/no. To  
170 determine what type of work the women considered physically demanding the following  
171 additional questions were asked: “How long do you stand or walk for in your occupation?”, with



172 the chosen cut-off of more than 50% of their working hours , and “How often do you experience  
173 heavy lifting at your workplace?”, with the chosen cut-off of more than 10-20 times per day. The  
174 data on vacuum and forceps deliveries was collected from electronic medical records (PARTUS)  
175 and pooled to ensure adequate numbers to fit the regression model, and included all deliveries  
176 were vacuum/forceps was used. SUI and POP at mid-pregnancy were highly correlated with the  
177 same symptoms at 12 months postpartum and thus not included in the regression analysis  
178 ( $p < 0.001$ ). The World Health Organization (WHO) classification was used to categorize body  
179 mass index (BMI) at 12 months following delivery: underweight  $\leq 18.4$ , normal weight 18.5-24.9,  
180 overweight 25-29.9, and obese  $> 30$  (27). Subgroup analyses, using the Chi-Square test and  
181 independent sample t-test, were undertaken on the data of women who exercised from 6 weeks to  
182 12 months postpartum compared to those who did not continue exercising until 12 months  
183 postpartum. Furthermore, we compared those not exercising either 6 weeks- or 12 months  
184 postpartum and those only exercising 12 months postpartum with those who had commenced  
185 exercising at 12 months postpartum. A p-value of  $\leq 0.05$  was considered statistically significant.  
186 No power calculation was undertaken for this study.

187

188

189

190

191

192

193 RESULTS

194 Flow of participants

195 At 6 weeks postpartum, 281 primiparous women remained enrolled in the study. Eighty-six  
196 (31%) women were classified as Exercisers ( $\geq 3$  times  $\geq 30$  min/week) and 195 (69%) as Non-  
197 exercisers. No differences on background variables at 6 weeks postpartum were found between  
198 the two groups (Table 1). Of those classified as Exercisers 6 weeks postpartum, 14 (5%) reported  
199 participating in high-impact exercise involving running and jumping  $\geq 3$  times  $\geq 30$  min/week.  
200 None of the high-impact exercisers were observed as statistical outliers in the univariate analysis,  
201 as assessed by normal Q-Q plots. This subgroup of high-impact exercisers were no different from  
202 the total sample of Exercisers with respect to VRP, PFM strength or PFM endurance (p-value:  
203 0.16-0.94) or symptoms of SUI and POP (p-value: 0.66-0.98). Thus, these women were included  
204 in the Exercisers group for all analysis.

205 Thirty-five (12%) were lost to follow up and 88 (29%) were excluded during the recruitment  
206 phase (Figure 1). The reason most participants were excluded was because they were  
207 participating in an RCT looking at the effects of PFMT.

208 We then compared women lost to follow up or excluded to the women still participating in the  
209 study at 12 months postpartum (n=177) with regard to various background variables. The only  
210 difference between the groups was that women still participating in our study at 12 months  
211 postpartum, reported undertaking PFMT  $\geq 3$  times per week compared to those lost to follow up  
212 or excluded 74 (42%) versus 29 (28%) (p-value 0.03), respectively.

213 How does general exercise 6 weeks postpartum impact on PFM variables and symptoms of SUI  
214 and POP 12 months postpartum?

215 One hundred and seventy-seven primiparous women were assessed at 12 months postpartum  
216 (mean 13 months, SD 0.8). Table 2 presents the mean difference with SD between Exercisers  
217 (n=57) and Non-Exercisers (n=120) on VRP, PFM strength and PFM endurance measured at 6  
218 weeks and 12 months postpartum. Table 3 presents numbers and percentages of symptoms of  
219 SUI and POP measured at 12 months postpartum for both groups. There were no statistically  
220 significant relationships found between Exercisers and Non-Exercisers on any variables (Table 2  
221 and 3).

222 In the adjusted model for Table 3, women with BMI between 25-29.9 were twice as likely to  
223 report SUI (OR=2.2, 95%CI: 1.0, 4.7) and women with BMI >30 were over three times more  
224 likely to report SUI (OR=3.3, 95%CI: 1.2, 9.4) at 12 months postpartum. Women with  
225 physically strenuous occupations (88/177 (50%)) were over three times more likely to report POP  
226 at 12 months postpartum (OR=3.0, 95%CI: 1.2, 7.3). In response to being asked what type of  
227 work participants considered physically demanding, 79 of 177 (45%) reported walking and/or  
228 standing more than 50% of the working day and 16 of 177 (9%) reported daily heavy lifting more  
229 than 10-20 times a day.

230 The subgroup analysis of maintenance of exercise from 6 weeks to 12 months postpartum  
231 showed no differences in PFM variables or symptoms of SUI and POP between those continuing  
232 exercise from 6 weeks to 12 months postpartum (n=34) compared to: those who did not continue  
233 to exercise until 12 months postpartum (n=52), those not exercising at either time-point  
234 postpartum (n=50), or those starting exercise 12 months postpartum (n=41) (p=0.32-0.95).

235

236 DISCUSSION

237 In the present study, regular general exercise performed at 6 weeks postpartum did not negatively  
238 influence the PFM or increase the risk of symptoms of SUI or POP 12 months postpartum. Our  
239 definition of Exercisers included many participants who were regularly participating in only low-  
240 impact types of activities. Consequently, we cannot extrapolate our findings to women  
241 participating in high-impact exercise. Future studies are warranted to establish the effect of more  
242 strenuous exercise on the pelvic floor in the early postpartum period (for example: running,  
243 repetitive jumping and heavy weightlifting) as women participating in such activities may be at  
244 increased risk (2,26). UI during high-impact exercise is common (10) and high-impact exercise  
245 prior to pregnancy has been found to increase UI in primiparous women 12 months postpartum  
246 (28).

247 Unfortunately, 52 out of 120 (43%) of the women classified as Exercisers 6 weeks postpartum  
248 were no longer exercising regularly 12 months following childbirth. In a recent survey by Dakic  
249 et al (29), SUI was identified as a barrier for discontinuing exercise. However, in our subgroup  
250 analyses we found no differences in PFM variables or symptoms of SUI or POP between those  
251 women who continued with regular exercise from 6 weeks to 12 months postpartum versus those  
252 women who ceased undertaking regular exercise during this period.

253 Little research appears to have been undertaken into the effects of general exercise on PFM  
254 function or PFD after childbirth. A recent study with data from the same cohort as our study  
255 found that those who were exercising regularly at gestational week 37, had stronger PFMs than  
256 Non-exercisers (30). The results of this study, however, are not comparable with that from Bø et  
257 al (30) as our participants were tested postpartum.

258 We found that overweight and obesity was a risk factor for SUI at 12 months postpartum.  
259 Furthermore, those women with physically strenuous occupations experienced more symptoms of  
260 POP at 12 months postpartum. Physically strenuous jobs are a known risk factor for PFD in the  
261 general female population (26). Forty-five percent of the women in our study reported standing  
262 and/or walking more than 50% of their working hours, and 9% reported heavy lifting at work on  
263 a daily basis. Our questionnaire did not define “heavy lifting”. Furthermore, we are not aware of  
264 any cut-off values that defines “how much standing and/or walking” or “how much heavy lifting”  
265 is considered harmful. This may have biased responses. Type of physically strenuous occupations  
266 or tasks in relation to risk of symptoms of POP warrants further investigation. Another finding  
267 was that symptoms of SUI and POP at mid-pregnancy were highly associated with the same  
268 symptoms at 12 months postpartum. This highlights the importance of addressing these issues  
269 prior to delivery. BMI is a potentially modifiable risk factor. The consequence of strenuous  
270 exercise on PFD in women with high BMI postpartum is not known and requires further research.  
271 Previous research from our study group found that the prevalence of anatomic POP was low  
272 throughout the study period (0-9%), while vaginal bulge symptoms were more prevalent (16-  
273 23%) (8). This may be explained by the ultrasound examination of anatomic POP being  
274 undertaken in supine, but symptoms of POP being experienced with many activities of daily  
275 living in an upright or standing position. The findings regarding POP is therefore a point of  
276 consideration.

277 The strengths of our study were the collection of data on general exercise training in the early  
278 postpartum period and the influence on symptoms of SUI, POP and AI at 12 months postpartum.  
279 To date, there is limited knowledge on how to guide women on postpartum exercise as we have  
280 been unsure of its effect on pelvic floor function. We chose to study the outcomes at 12 months

281 postpartum to minimize the effect of childbirth on the pelvic floor (3). Another strength was that  
282 few women were lost to follow up. Notably, the majority of the 88 (29%) who were excluded  
283 from the study were excluded because they were participating in an RCT looking at the effects of  
284 PFMT. Inclusion of these participants would have affected the outcomes of this study given it is  
285 well established that PFMT has level 1A scientific evidence for the treatment of UI and POP  
286 (31,32).

287 Due to the finding that women who remained in the study until 12 months postpartum undertook  
288 more PFMT at 6 weeks postpartum than those who were lost to follow up or excluded, we used  
289 PFMT as a variable in the regression analysis. Exercising the PFM  $\geq$  three times per week did not  
290 influence PFM function or the presence of SUI or POP. Our findings that 43% did not continue  
291 with regular general exercise between 6 weeks and 12months postpartum, highlights that reasons  
292 for ceasing exercise during the first 12months after delivery should be investigated in the future.

293 A limitation of our study was that there was a large variation in the type of exercise being  
294 undertaken by the women. We believe, however, that this information is still valuable as it  
295 provides information on what types of exercise women like to participate in during the early  
296 postpartum period.

297 The results from our study shows that starting regular general exercise within the first 6 weeks  
298 postpartum does not negatively influence the PFM, which is an important message to impart to  
299 postpartum women and health care providers. Although there has been concern that high-impact  
300 exercise may be harmful to the PFM (2,26,32), the subgroup analyses of the 14 women in our  
301 study who undertook high-impact exercise  $\geq 3$  times  $\geq 30$  min/week showed no difference in PFM  
302 function or the presence of SUI or POP as compared to the total sample. However, due to the  
303 small numbers we cannot rule out the risk of a type II-error. We acknowledge the limitation that

304 the ICIQ-VS and the ICIQ-B have not yet been validated in the Norwegian language. However,  
305 the questions on these outcome measures are straight forward and used worldwide, and it was  
306 important to investigate the effects of general exercise on POP and AI in the early postnatal  
307 period (21,23,24). A limitation was that there was no power calculation undertaken prior to the  
308 study. To be able to study the impact of exercise on AI, and in addition the impact of type of  
309 exercise (especially strenuous exercises) on the pelvic floor, studies with a larger sample sizes are  
310 needed in this population. Further limitations were that the women were asked to recall both  
311 presence of PFD and frequency of exercise, which may have biased the results. Participants were  
312 asked to recall whether or not they have been exercising the last four weeks. Consequently, we do  
313 not know the exact time of start of exercise. Our data may have been more precise had we used a  
314 shorter time interval. This questionnaire, however, has been widely used in previous studies,  
315 including the Norwegian Mother and Child Cohort (33), and is considered to be representative  
316 for this population. Frequency of exercise was also based on self-report. Objective measures of  
317 physical activity and exercise were not collected as this study was a part of a larger cohort study  
318 on pelvic floor changes throughout pregnancy and postpartum (3). Furthermore, our study cannot  
319 be generalize to other ethnic groups.

320

321

322

323

324

325

326 ACKNOWLEDGEMENTS:

327 The authors would like to thank all the women who participated in this project. We would also  
328 like to extend our gratitude to the staff working at the Department of Obstetrics and Gynaecology  
329 at Akershus University Hospital, Lørenskog, Norway especially: midwife Tone Breines  
330 Simonsen for recruiting participants, gynecologist Franziska Siafarikas and Jette Stær-Jensen,  
331 and physical therapists Gunvor Hilde and Kristin Gjestland for clinical testing and data entry. We  
332 thank Natalie Michelle Doig Evensen for English grammar editing.

333 SOURCES OF FUNDING:

334 The present study was part of a larger cohort study which was funded by the the South-Eastern  
335 Norway Regional Health Authority and the Norwegian Research Council. Extra funding for this  
336 specific study was granted by EXTRA funds from the Norwegian Foundation for Health and  
337 Rehabilitation and the Norwegian Women's Public Health Association.

338 CONFLICT OF INTEREST: None

339

340

341

342

343

344

345



346 REFERENCES

- 347 1. Gynecologists. TACoOa. ACOG Committee opinion no. 650. Summary: physical activity and  
 348 exercise during pregnancy and the postpartum period. *Obstetrics & Gynecology*. 2015; 126(1326-  
 349 7.
- 350 2. Bo K, Artal R, Barakat R, et al. Exercise and pregnancy in recreational and elite athletes: 2016/17  
 351 evidence summary from the IOC Expert Group Meeting, Lausanne. Part 3-exercise in the  
 352 postpartum period. *Br J Sports Med*. 2017; 51(21): 1516-25.
- 353 3. Staer-Jensen J, Siafarikas F, Hilde G, Benth JS, Bo K, Engh ME. Postpartum recovery of levator  
 354 hiatus and bladder neck mobility in relation to pregnancy. *Obstetrics & Gynecology*. 2015;  
 355 125(3): 531-9.
- 356 4. Reimers C, Staer-Jensen J, Siafarikas F, Saltyte-Benth J, Bo K, Ellstrom Engh M. Change in pelvic  
 357 organ support during pregnancy and the first year postpartum: a longitudinal study. *British*  
 358 *Journal of Obstetrics & Gynaecology*. 2016; 123(5): 821-9.
- 359 5. Sievert KD AB, Toomey PA, Robinson D, Milsom I, Koelbl H, Abrams P, Cardozo L, Wein A, Smith  
 360 AL, Newman DK. Can we prevent incontinence?: ICI-RS 2011. *Neurourology & Urodynamics*.  
 361 2012; 31(390-9).
- 362 6. Thom DH, Rortveit G. Prevalence of postpartum urinary incontinence: a systematic review. *Acta*  
 363 *Obstet Gynecol Scand*. 2010; 89(12): 1511-22.
- 364 7. Johannessen HH, Wibe A, Stordahl A, Sandvik L, Backe B, Morkved S. Prevalence and predictors  
 365 of anal incontinence during pregnancy and 1 year after delivery: a prospective cohort study.  
 366 *British Journal of Obstetrics & Gynaecology*. 2014; 121(3): 269-79.
- 367 8. Reimers C, Staer-Jensen JE, Siafarikas F, Bo K, Engh ME. Association between vaginal bulge and  
 368 anatomical pelvic organ prolapse during pregnancy and postpartum: an observational study. *Int*  
 369 *Urogynecol J*. 2018; 29(3): 441-8.
- 370 9. Bø K. Urinary incontinence, pelvic floor dysfunction, exercise and sport. *Sports Medicine*. 2004;  
 371 34(7): 451-64.
- 372 10. Nygaard I, Shaw JM. Physical activity and the pelvic floor. *Am J Obstet Gynecol*. 2016; 214(2):  
 373 164-71.
- 374 11. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies  
 375 in Epidemiology (STROBE) statement: guidelines for reporting observational studies *J Clin*  
 376 *Epidemiol*. 2008; 61(4): 344-9.
- 377 12. Bouchard C, Shephard R, J. Physical Activity, Fitness , and Health: The Model and Key Concepts.  
 378 In: Bouchard C, Shephard R, J. , Stephens R, eds. Physical Activity, Fitness, and Health. Consensus  
 379 Statement. Champaign: Human Kinetics Publishers; 1993. p. 11-24.
- 380 13. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness:  
 381 definitions and distinctions for health-related research. *Public Health Rep*. 1985; 100(126-31).
- 382 14. Garber CE, Blissmer B, Deschenes MR, et al. ACSM Position Stand: Quantity and Quality of  
 383 Exercise for Developing and Maintaining Cardiorespiratory, Musculoskeletal, and Neuromotor  
 384 Fitness in Apparently Healthy Adults: Guidance for Prescribing Exercise. *Medicine & Science in*  
 385 *Sports & Exercise*. 2011: 1334-59.
- 386 15. Kegel AH. Stress incontinence and genital relaxation; a nonsurgical method of increasing the  
 387 tone of sphincters and their supporting structures. *Ciba Clin Symp*. 1952; 4(2): 35-51.
- 388 16. Bø K. Pressure measurements during pelvic floor muscle contractions: the effect of different  
 389 positions of the vaginal measuring device. *Neurourology & Urodynamics*. 1992; 11(2): 107-13.
- 390 17. Bø K, Hagen RR, Kvarstein B, Larsen S. Pelvic floor muscle exercise for the treatment of female  
 391 stress urinary incontinence. II. Validity of vaginal pressure measurements of pelvic muscle

- 392 strength and the necessity of supplementary methods for control of correct contraction.  
393 *Neurourology & Urodynamics*. 1990; 9(5): 479-87.
- 394 18. Bø K, Hagen R, Kvarstein B, Larsen S. Pelvic floor muscle exercise for the treatment of female  
395 stress urinary incontinence: I. Reliability of vaginal pressure measurements of pelvic muscle  
396 strength. *Neurourology & Urodynamics*. 1990; 9(5): 471-7.
- 397 19. Tennfjord MK, Engh ME, Bø K. An intra- and interrater reliability and agreement study of vaginal  
398 resting pressure, pelvic floor muscle strength, and muscular endurance using a manometer. *Int*  
399 *Urogynecol J*. 2017; 28(10): 1507-14.
- 400 20. Wilmore JH, Costill DL. *Physiology of sport and exercise*. 2 ed. Leeds: Human Kinetics; 1999.
- 401 21. Avery K, Donovan J, Peters TJ, Shaw C, Gotoh M, Abrams P. ICIQ: a brief and robust measure for  
402 evaluating the symptoms and impact of urinary incontinence. *Neurourology & Urodynamics*.  
403 2004; 23(4): 322-30.
- 404 22. Abrams P, Avery K, Gardener N, Donovan J, Board IA. The International Consultation on  
405 Incontinence Modular Questionnaire: [www.iciq.net](http://www.iciq.net). *J Urol*. 2006; 175(3 Pt 1): 1063-6; discussion  
406 6.
- 407 23. Price N, Jackson S, Avery K, Brookes ST, Abrams P. Development and psychometric evaluation of  
408 the ICIQ Vaginal Symptoms Questionnaire: the ICIQ-VS. *British Journal of Obstetrics &*  
409 *Gynaecology*. 2006; 113(700-12).
- 410 24. Cotterill N, Norton C, Avery KN, Abrams P, Donovan JL. Psychometric evaluation of a new  
411 patient-completed questionnaire for evaluating anal incontinence symptoms and impact on  
412 quality of life: the ICIQ-B. *Diseases of the Colon & Rectum*. 2011; 54(10): 1235-50.
- 413 25. Sultan AH, Monga A, Lee J, et al. An International Urogynecological Association  
414 (IUGA)/International Continence Society (ICS) joint report on the terminology for female  
415 anorectal dysfunction. *Int Urogynecol J*. 2017; 28(1): 5-31.
- 416 26. Milson I, Altman D, Cartwright R, et al. Epidemiology of Urinary Incontinence (UI) and Other  
417 Lower Urinary Tract Symptoms (LUTS), Pelvic Organ Prolapse (POP) and Anal Incontinence (AI).  
418 In: Abrams P, Cardozo L, Khoury S, Wein A, eds. Incontinence. 5th ed.: ICUD Consultations; 2013.  
419 p. 15-108.
- 420 27. Europe WHORof. Body mass index - BMI. 2019; [http://www.euro.who.int/en/health-](http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi)  
421 [topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi](http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi). Accessed 02.12,  
422 2019.
- 423 28. Durnea CM, Khashan AS, Kenny LC, et al. What is to blame for postnatal pelvic floor dysfunction  
424 in primiparous women-Pre-pregnancy or intrapartum risk factors? *European Journal of*  
425 *Obstetrics, Gynecology, & Reproductive Biology*. 2017; 214(36-43).
- 426 29. Dakic J, Cook J, Lin K, Hay-Smith J, Frawley H. The impact of pelvic floor dysfunction on exercise.  
427 In: *International Continence Society*; Year 05.09.2019: Sweden.
- 428 30. Bø K, Engh ME, Hilde G. Regular exercisers have stronger pelvic floor muscles than nonregular  
429 exercisers at midpregnancy. *American Journal of Obstetrics & Gynecology*. 2018; 218(427): 1-5.
- 430 31. Caccaiari LP, Dumoulin C, Hay-Smith EJ. Pelvic floor muscle training versus no treatment, or  
431 inactive control treatments, for urinary incontinence in women: a cochrane systematic review  
432 abridged republication. *Braz J Phys Ther*. 2019; 23(2): 93-107.
- 433 32. Dumoulin C, Adewuyi t, Booth J, et al. Adult conservative management. In: Abrams P, Cardozo L,  
434 Wagg A, Wein A, eds., eds. Incontinence 6th Edition (2017). Bristol UK: International Continence  
435 Society; 2017. p. 1443-628.
- 436 33. Owe KM, Nystad W, Bø K. Correlates of regular exercise during pregnancy: the Norwegian  
437 Mother and Child Cohort Study. *Scandinavian Journal of Medicine & Science in Sports*. 2009;  
438 19(5): 637-45.

439

440 Table 1. Background variables of primiparous women 6 weeks postpartum and differences between Exercisers and Non-exercisers.

	Total study population n=281	Exercisers n=86	Non- exercisers n=195	Difference between Exercisers/ Non- exercisers
Age: (years)	29 (SD 4.3)	29 (SD 4.1)	29 (SD 4.4)	95%CI: (-1.36,0.82)
BMI: (kg/m <sup>2</sup> )	26 (SD 3.9)	26 (SD 3.9)	26 (SD 3.9)	95%CI: (-1.33, 0.66)
Married/cohabitant	269 (96%)	81 (94%)	188 (96%)	0.60
College/university degree	213 (76%)	62 (72%)	151 (77%)	0.42
Smoking	16 (5.7%)	3 (3.5%)	13 (6.7%)	0.44
Breastfeeding	265 (94%)	82 (95%)	183 (94%)	0.83

Pelvic floor muscle training $\geq 3$ times per week	103 (37%)	31 (36%)	72 (37%)	1.0
Normal vaginal delivery	193 (69%)	61 (71%)	132 (68%)	0.48
Vacuum/forceps*	46 (16%)	10 (12%)	36 (19%)	0.21
Pre-labor CS (cervix $< 3$ cm)	20 (7.1%)	8 (9.3%)	12 (6.2%)	0.48
Intrapartum CS (cervix $\geq 3$ cm)	22 (7.8%)	7 (8.1%)	15 (7.7%)	0.48
Prolonged second stage (n=231)	39 (16%)	10 (14%)	29 (18%)	0.58

Pelvic organ prolapse	57 (20%)	16 (19%)	41 (21%)	0.76
Stress urinary incontinence	79 (28%)	20 (23%)	59 (30%)	0.29
Anal incontinence	4 (1.4%)	1 (1.20%)	3 (1.5%)	1.0
Pelvic floor muscle variables:				
Vaginal resting pressure (cm H <sub>2</sub> O)	32 (SD 9.0)	32 (SD 8.5)	32 (SD 9.2)	95%CI: (-2.4, 2.1)
Pelvic floor muscle strength (cm H <sub>2</sub> O)	19 (SD 14)	20 (SD 14)	18 (SD 14)	95%CI: (-5.7, 1.5)
Pelvic floor muscle endurance (cm H <sub>2</sub> O sec)	131 (SD 105)	145 (SD 109)	125 (SD 103)	95%CI: (-47, 6.1)

441 BMI= body mass index, CS=caesarean section. Continuous variables are given as means with standard deviation (SD) and categorical  
442 variables are given as frequencies (n) with percentages (%). The difference between Exercisers and Non-exercisers is given as 95% CI  
443 or p-value. \* Forty-four vacuum -and two forceps deliveries.

444

445

446 Table 2. Associations between Exercisers and Non-exercisers at 6 weeks postpartum and pelvic floor muscle variables 12 months  
 447 postpartum.

Dependent variable	Exercise	Mean (SD)	Unadjusted B (95%CI)	P-value	Adjusted B (95%CI)*	P-value*
Vaginal resting pressure (cm H <sub>2</sub> O)	Exercisers (n=57)	35 cm H <sub>2</sub> O (SD 7.8)				
	Non-exercisers (n=120)	35 cm H <sub>2</sub> O (SD 9.3)	-0.04 (-3.4, 2.1)	0.64	-0.02 (-3.2, 2.8)	0.75
PFM strength (cm H <sub>2</sub> O)	Exercisers (n=57)	36 cm H <sub>2</sub> O (SD 17)				
	Non-exercisers (n=120)	34 cm H <sub>2</sub> O (SD 20)	0.03 (-4.7, 7.4)	0.67	0.02 (-5.2, 6.8)	0.78

PFM endurance (cm H <sub>2</sub> O sec)	Exercisers (n=57)	272 cm H <sub>2</sub> O sec (SD 141)				
	Non-exercisers (n=120)	278 cm H <sub>2</sub> O sec (SD 178)	-0.02 (-59, 46)	0.81	-0.03 (-63, 41)	0.66

448 Differences between Exercisers and Non-exercisers at 6 weeks postpartum on VRP, PFM strength and PFM endurance were assessed  
449 using the independent sample t-test and presented as means with SD. Standard multiple linear regressions with associations presented  
450 as standardised beta coefficients (B) with 95% confidence intervals (CI) and p-values. PFM=pelvic floor muscles. Adjusted for age,  
451 physically strenuous occupations, body mass index (kilograms per square meter), pelvic floor muscle training  $\geq 3$  times per week,  
452 vacuum/forceps. \*n=176

453

454

455

456

457

458 Table 3. Associations between Exercisers and Non-exercisers at 6 weeks postpartum and stress urinary incontinence and pelvic organ  
 459 prolapse 12 months postpartum.

Dependent variable	Exercise	Frequencies (n) (%)	OR unadjusted (95% CI)	Unadjusted p-value	OR adjusted (95% CI)*	Adjusted p-value*
SUI n=57	Exercisers (n=57)	13 (22%)				
	Non-exercisers (n=120)	44 (37%)	0.51 (0.25, 1.1)	0.07	0.53 (0.25, 1.1)	0.10
POP n=33	Exercisers (n=57)	8 (14%)				
	Non-exercisers (n=120)	25 (21%)	0.62 (0.26, 1.5)	0.28	0.57 (0.23, 1.4)	0.22

460 SUI=stress urinary incontinence, POP=pelvic organ prolapse. Adjusted for age, physically strenuous occupations, body mass index  
 461 (kilograms per square meter), pelvic floor muscle training  $\geq 3$  times per week, vacuum/forceps. \*n=176



462 Differences between Exercisers and Non-exercisers on symptoms of SUI and POP were assessed using the Chi-Square test and  
463 presented with frequencies (n) and percentages. Logistic regression model with associations presented with odds ratio (OR) with 95%  
464 confidence intervals (CI) and p-values were undertaken.

465 Figure 1. Flowchart of the study participants included in the study at 12 months postpartum.

466 Boxes on the left represent the actual number of participants in the study at all times. Boxes

467 on the right represent participants lost to follow up, excluded or those with missing data.

468

469

470

471

472

473

474

475

476

477

478

479


480

481

482

483

484 Figure 1. Flowchart of the study participants included in the study at 12 months postpartum.



\*Missing at one timepoint is no exclusion but indicates that the participant may participate in the study at the next timepoint.